



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

B. poliocephala (Reichenow), *B. xavieri* (Oust.), *B. notata* (Cass.), *B. eximia* (Hartl.), and *B. canicapilla* (Hartl.).—CHAS. W. RICHMOND, Washington, D. C.

Barn Swallows (*Hirundo erythrogastra*).—Within a few yards of the house occupied by Mr. John R. Sandsbury during the time he is caring for the Terns on Muskeget Island, and where I make my headquarters when visiting there, is an old shed or boathouse which has several apertures. This shed has been used as a nesting place for the past six years by apparently the same pair of Barn Swallows. At my request Mr. S. made a few notes on these birds, which arrived this year (1898) on May 29. It is their custom to repair the old nest, they never having built any since the first one. Four young birds were hatched this season. The old birds would occasionally fly into the sitting-room of the house, but were always frightened on getting inside. When I was visiting Muskeget this summer (July 2-5, 1898), I found, in addition to the old pair of birds, still another pair, apparently birds of last year, assisting in feeding the four young ones in the nest. This they continued to do up to July 10, the date on which the young left the nest. On this date they were all flying about together, the young going at intervals to the nest to rest. On July 11, there were only the two original old birds and the four young ones, and they remained around until July 19, the young returning to the nest every night. The young birds were so tame that they would alight on, and even run over Mr. Sandsbury's fingers while he rested his hand upon a beam which was near the nest. They returned occasionally up to August 1, but were not so tame, alighting on top of the shed and on the clothes line near the house. At this latter date the group consisted of the two old birds and the four young.—GEORGE H. MACKAY, Nantucket, Mass.

Another Example of Curious Nesting of the American Redstart.—Mr. Verdi Burtch, in the October Auk, 1898, recorded a curious example of the American Redstart's nesting, and having had a somewhat similar experience, it may be of interest to record it.

June 3, 1898, I had been collecting about a swamp in the vicinity of Dorchester, Mass., and at noon sought the shade of a wood lot near. A female Redstart (*Setophaga ruticilla*) at once attracted my attention by her queer ways. I retired for a short distance and the bird settled upon a Vireo's nest, which was situated four and a half feet above the ground in a sapling. It contained five Redstart eggs. One of these was entirely buried beneath the others, in a thick lining of horse hair. The yolk of this egg had settled and hardened. The other four were fresh.

As numbers of Redstarts' eggs are annually stolen by boys from this wood, it may be possible that the following theory accounts for this strange thing. An incomplete set of Redstart eggs was taken; the female laid in the Vireo nest during the absence of the owners rather than deposit